







2.

$$f(x) = x^3 + 2x^2 - 3x - 11$$

(a) Show that  $f(x) = 0$  can be rearranged as

$$x = \sqrt{\left(\frac{3x+11}{x+2}\right)}, \quad x \neq -2.$$

(2)

The equation  $f(x) = 0$  has one positive root  $\alpha$ .

The iterative formula  $x_{n+1} = \sqrt{\left(\frac{3x_n+11}{x_n+2}\right)}$  is used to find an approximation to  $\alpha$ .

(b) Taking  $x_1 = 0$ , find, to 3 decimal places, the values of  $x_2$ ,  $x_3$  and  $x_4$ .

(3)

(c) Show that  $\alpha = 2.057$  correct to 3 decimal places.

(3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

















Question 4 continued

Leave  
blank

Lined area for writing the answer to Question 4.

(Total 9 marks)

Q4



5. Sketch the graph of  $y = \ln|x|$ , stating the coordinates of any points of intersection with the axes.

(3)

Leave  
blank



**Question 5 continued**

Leave  
blank

**(Total 3 marks)**

**Q5**

13

**Turn over**



6.




Figure 1

Figure 1 shows a sketch of the graph of  $y = f(x)$ .

The graph intersects the  $y$ -axis at the point  $(0, 1)$  and the point  $A(2, 3)$  is the maximum turning point.

Sketch, on separate axes, the graphs of

- (i)  $y = f(-x) + 1$ ,
- (ii)  $y = f(x + 2) + 3$ ,
- (iii)  $y = 2f(2x)$ .

On each sketch, show the coordinates of the point at which your graph intersects the  $y$ -axis and the coordinates of the point to which  $A$  is transformed.

(9)



**Question 6 continued**

Leave  
blank



**Question 6 continued**

Leave  
blank





Question 6 continued

Leave  
blank

(Total 9 marks)

Q6





Leave blank

7. (a) By writing  $\sec x$  as  $\frac{1}{\cos x}$ , show that  $\frac{d(\sec x)}{dx} = \sec x \tan x$ .

(3)

Given that  $y = e^{2x} \sec 3x$ ,

(b) find  $\frac{dy}{dx}$ .

(4)

The curve with equation  $y = e^{2x} \sec 3x$ ,  $-\frac{\pi}{6} < x < \frac{\pi}{6}$ , has a minimum turning point at  $(a, b)$ .

(c) Find the values of the constants  $a$  and  $b$ , giving your answers to 3 significant figures.

(4)

Lined area for student answers.









8. Solve

$$\operatorname{cosec}^2 2x - \cot 2x = 1$$

for  $0 \leq x \leq 180^\circ$ .

(7)

Leave  
blank





Leave  
blank

**Question 8 continued**

Lined writing area for the answer.

**Q8**

**(Total 7 marks)**









**Question 9 continued**

Leave  
blank

Lined area for writing the answer to Question 9.





