Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate No.			6	6	7	4	/	0	1	Signature	

Paper Reference(s)

6674/01

Edexcel GCE

Further Pure Mathematics FP1 Advanced/Advanced Subsidiary

Monday 19 June 2006 – Morning

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Green)

Items included with question papers
Nil

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initial(s) and signature. Check that you have the correct question paper.

You must write your answer for each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the examiner. Answers without working may gain no credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

©2006 Edexcel Limited

N20461A W850/R6674/57570 3/3/3/3/10,900





Turn over

Total

Examiner's use only

Team Leader's use only

Question Number

2

3

5

6

7

8



$2z + iw = -1,$ $z - w = 3 + 3i.$ (a) Use algebra to find z, giving your answer in the form $a + ib$, where a and b (b) Calculate $\arg z$, giving your answer in radians to 2 decimal places.	are real.
(a) Use algebra to find z , giving your answer in the form $a + ib$, where a and b	
(b) Calculate $\arg z$, giving your answer in radians to 2 decimal places.	
	(2)

Question 1 continued	
Question 1 continued	




Figure 1 shows a curve C with polar equation $r=4a\cos 2\theta$, $0\leqslant \theta\leqslant \frac{\pi}{4}$, and a line m with polar equation $\theta=\frac{\pi}{8}$. The shaded region, shown in Figure 1, is bounded by C and m. Use calculus to show that the area of the shaded region is $\frac{1}{2}a^2(\pi-2)$.

Question 2 continued	

Given that $3x \sin 2x$ is a particular integral of the differential equation

$$\frac{\mathrm{d}^2 y}{\mathrm{d}x^2} + 4y = k\cos 2x \;,$$

where k is a constant,

(a) calculate the value of k,

(4)

(b) find the particular solution of the differential equation for which at x = 0, y = 2, and for which at $x = \frac{\pi}{4}$, $y = \frac{\pi}{2}$.

(4)

Duration 2 continued	
Question 3 continued	

 x⁴-6x³+19x²-36x+78=0, (a) solve the equation completely, (b) show on a single Argand diagram the four points that represent the roo equation. 	(7)
(b) show on a single Argand diagram the four points that represent the roo	
(b) show on a single Argand diagram the four points that represent the roo	
(b) show on a single Argand diagram the four points that represent the roo equation.	ts of the
equation.	
	(2)

	Leave blank
Question 4 continued	
	Q4
(Total 9 marks)	

5. Given that for all real values of r,

$$(2r+1)^3-(2r-1)^3=Ar^2+B$$
,

where A and B are constants,

(a) find the value of A and the value of B.

(2)

(b) Hence, or otherwise, prove that $\sum_{r=1}^{n} r^2 = \frac{1}{6} n(n+1)(2n+1).$

(5)

(c) Calculate $\sum_{r=1}^{40} (3r-1)^2$.

(3)

10

	Le bla
Question 5 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_

Question 5 continue		
		_
		-
		—
		-

Question 5 continued	Lo bl

	1
Leave	
blank	

6.	$f(x) = 0.25x - 2 + 4 \sin \sqrt{x}.$	
	(a) Show that the equation $f(x) = 0$ has a root α between $x = 0.24$ and $x = 0.28$.	
		(2)
	(b) Starting with the interval [0.24, 0.28], use interval bisection three times to find interval of width 0.005 which contains α .	an
		(3)
	The equation $f(x) = 0$ also has a root β between $x = 10.75$ and $x = 11.25$.	
	(c) Taking 11 as a first approximation to β , use the Newton-Raphson process on $f(x)$ or	nce
	to obtain a second approximation to β . Give your answer to 2 decimal places.	(6)
		(0)

	L	_eave
Question 6 continued		

	L
Question 6 continued	
	(Total 11 marks)

7.	(a) Use algebra to find the exact solutions of the equation			
		$ 2x^2 + x - 6 = 6 - 3x.$		

(b) On the same diagram, sketch the curve with equation $y = |2x^2 + x - 6|$ and the line with equation y = 6 - 3x.

(3)

(6)

(c) Find the set of values of x for which

$$|2x^2+x-6| > 6-3x$$
.

(3)

18

	Lea bla
Question 7 continued	

Question 7 continued			
			_
			_
			-
			_
			_
			-
			_
			_
			_
			-
			-
			_
			_
			-
			_
			-
			_
			_
			_
			_
			_

Question 7 continued	
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-
	-

L	eave	
h	onla	

8. During an industrial process, the mass of salt, $S \log t$, dissolved in a liquid t minutes after the process begins is modelled by the differential equation $\frac{dS}{dt} + \frac{2S}{120 - t} = \frac{1}{4}, \qquad 0 \le t < 120.$

Given that S = 6 when t = 0,

(a) find S in terms of t,

(8)

(b) calculate the maximum mass of salt that the model predicts will be dissolved in the liquid at any one time during the process.

(4)

Question 8 continued	Leav blan

Question 8 continued	
	(Total 12 marks)
	TOTAL FOR PAPER: 75 MARKS